

Suggestions for skills assessment with the **LIFEPAK® 35** Monitor/Defibrillator

Pediatric features skills assessment

Name:Reviewer:				
This skills checklist assesses a trained user's ability to care for the pediatric patient with the LIFEPAK 35 monitor/defibrillator. This limited checklist does not cover all information and skills required to operate the device safely and effectively. This checklist is designed to be completed after receiving or viewing in-service training by a qualified instructor. Refer to the LIFEPAK 35 monitor/defibrillator Operating Instructions for complete directions for use, indications, contraindications, warnings, eautions, and potential adverse events. CAPITAL LETTERS indicate labels, screen messages, and voice prompts. "Push" is used when referring to keypad buttons, and Press" is used when referring to softkeys on the touchscreen indicated by italics				
Assessment criteria	Complete	Incomplete	Comments	
1. Pushes the POWER button.			For this assessment, instruct the participant that they are caring for a 3-year-old 14kg patient.	
 2. Changes Patient Type: Press the ALARMS button at the bottom of the Home Screen. Press PEDIATRIC. Press anywhere else on the touchscreen to close the ALARMS menu. 			Ask the participant to verbalize their selection for a patient less than 1 month old.	
 3. Enables Alarms: Press the ALARMS button at the bottom of the touchscreen. Press ALL ON in the ALARMS area of the dialog box. 			If alarms are set up to be OFF, complete this step.	
 4. Applies appropriately sized monitoring accessories including: NIBP Cuff Oximeter sensor ECG monitoring electrodes 				
 5. Confirms the pediatric default high, low and No Breath alarm limits for each parameter. Press the color bar at the left edge of the parameter. Press the ALARMS button on the slide-out menu. Press SET ALARMS to open the alarms menu for the parameter. The alarms are turned on to pediatric default limits, and the alarm limits menu appears. 				
Pediatric AED Mode				
6. Pushes the ANALYZE button.				
7. Prepares patient's skin for therapy electrode pad placement.			See patient skin preparation for instructions.	
8. Connects the pediatric therapy electrode pads to the therapy cable. Confirms therapy cable is connected to the device.				
9. Applies the therapy electrode pads to the patient's chest in the anterior-lateral position.				
10 Confirms PEDIATRIC MODE				

11. Continues to follow the screen messages and voice prompts provided by the device.	CPR timer is in real or elapsed time at the top of the screen.
 12. SHOCK ADVISED: Provides a verbal "Clear" and confirms all clear. Pushes the (SHOCK) button to deliver energy to the patient. Follows the START CPR prompt after shock delivery. Acknowledges CPR prompt and timer will appear and continue per duration specified per medical direction local protocols or default settings in the device. Continue to follow the screen messages and voice prompts provided by the AED. Verbalizes energy delivered. 	 Pressing the Speed Dial cancels the charge or charge with disarm after 60 seconds. CPR timer is in real or elapsed time.
Pediatric Manual Mode Defibrillation	
13. Enters Manual Mode by pressing press MANUAL DEFIB in the lower-right	
14. Confirms desired pediatric energy is selected, or select the desired energy using the Up/Down arrows or dial on the touchscreen.	
15. Presses CHARGE.	
16. Provides a verbal "Clear" and confirms all clear.	
17. Confirms ECG rhythm requires defibrillation and confirms available energy.	
18. Presses the (SHOCK) button on the defibrillator.	
19.(Optional): Presses SPEED DIAL to cancel the charge.	
20. Starts CPR according to established protocol. Presses	Confirm that cprINSIGHT is toggled on and illuminated green.



Scan here for additional resources

This document is intended solely for the use of healthcare professionals. A healthcare professional must always rely on his or her own professional clinical judgment when deciding whether to use a particular product when treating a particular patient. Stryker does not dispense medical advice and recommends that healthcare professionals be trained in the use of any particular product before using it.

The information presented is intended to demonstrate Stryker's product offerings. A healthcare professional must always refer to operating instructions for complete directions for use indications, contraindications, warnings, cautions, and potential adverse events, before using any of Stryker's products. Products may not be available in all markets because product availability is subject to the regulatory and/or medical practices in individual markets. Please contact your representative if you have questions about the availability of Stryker's products in your area. Specifications subject to change without notice.

Stryker or its affiliated entities own, use, or have applied for the following trademarks or service marks: LIFEPAK, Stryker. All other trademarks are trademarks of their respective owners or holders.

The absence of a product, feature, or service name, or logo from this list does not constitute a waiver of Stryker's trademark or other intellectual property rights concerning that name or logo.

EC-L35-ASTL-1169282_REV-0_en_us © 2024 Stryker



Manufactured by:

Physio-Control, Inc. 11811 Willows Road NE Redmond, WA, 98052 U.S.A. Toll free 800 442 1142 stryker.com/emergencycare